arXiv:1403.5393v1 [math.GR] 21 Mar 2014

BI-COMMUTATIVE AG-GROUPOIDS

M. RASHAD^A, I. AHMA$D^{A,*}$, AND M. SHAH^B

Abstract. A groupoid satisfying the left invertive law: $ab \cdot c = cb \cdot a$ is called an AG-groupoid and is a generalization of commutative semigroups. We consider the concept of bi-commutativity in AG-groupoids and thus introduce left commutative AG-groupoids, right commutative AG-groupoids and bi-commutative AG-groupoids. We provide a method to test an arbitrary Cayley's table for these AG-groupoids and explore some of the general properties of these AG-groupoids.

1. introduction

An AG-groupoid S is in general a nonassociative groupoid that satisfies the left invertive law, $ab.c = cb.a$. S is called medial if it satisfies the medial property, $ab.cd = ac.bd$. it is easy to prove that every AG-groupoid is medial. S is called an AG-monoid if it contains the left identity element. Every AG-monoid is paramedial[10], i.e it satisfies the identity, $ab.cd = db.ca$. Some of the well known AG-groupoids with their identities are listed in Table 2. Recently many new classes of AG-groupoids have been introduced by various researchers[9, 11, 12, 13]. These new classes are studied in a variety of papers like for instance[4, 5, 6, 7, 14]. A groupoid G is called left (resp. right) commutative groupoid if G satisfies the identity $(ab)c = (ba)c$ (resp. $a(bc) = a(cb)) \forall a, b, c \in G$. In this article we extend the concept of these groupoids to introduce some new classes of AG-groupoids, that we call a left commutative AG-groupoid or shortly an LC-AG-groupoid, a right commutative AG-groupoid or an RC-AG-groupoid, and a bi-commutative AG-groupoid or BC-AG-groupoid. We use the GAP [2] software and the relevant data of [3] to enumerate our new classes of AG-groupoids up to order 6. The following table illustrates nonassociative enumeration of these classes up to order 6.

Key words and phrases. AG-groupoids, AG-test, nuclear square AG-groupoids, transitive groupoids, alternative AG-groupoids.

*Corresponding author.

Order	3	4	5	6
Nonassociative AG-groupoids	8	269	31467	40097003
LC-AG-groupoids	6	194	22276	34845724
RC-AG-groupoids	2	52	1800	170977
BC-AG-groupoid	2	47	1558	150977

Table 1. Enumeration of BC-AG-groupoids up to order 6.

An AG-groupoid is a groupoid satisfying the left invertive law,

$$ab \cdot c = cb \cdot a. \tag{1.1}$$

We investigate our newly introduced classes of AG-groupoids and find some relations of these classes with already known classes of AG-groupoids[8, 15, 16]. We list some of the already known classes of AG-groupoids with their defining identities that will be used in the rest of this article.

AG-groupoid	satisfying identity
Left nuclear square AG-groupoid	$a^2(bc) = (a^2b)c$
Middle nuclear square AG-groupoid	$(ab^2)c = a(b^2c)$
Right nuclear square AG-groupoid	$(ab)c^2 = a(bc^2)$
T^1-AG-groupoid	$ab = cd \Rightarrow ba = dc$
Medial AG-groupoid	$ab \cdot cd = ac \cdot bd$
Paramedial AG-groupoid	$ab \cdot cd = db \cdot ca$
Flexible -AG-groupoid	$ab \cdot a = a \cdot ba$
AG-3-band-AG-groupoid	$a(aa) = (aa)a = a$
Left alternative AG-groupoid	$aa \cdot b = a \cdot ab$
Self-dual AG-groupoid	$a(bc) = c(ba)$

Table. 2

2. Bi-commutative-AG-groupoids

Definition 1. *An AG-groupoid S is called —*
(a) — a left commutative AG-groupoid (LC-AG-groupoid) if $\forall a, b, c \in S$,

$$(ab)c = (ba)c \tag{2.1}$$

(b) — a right commutative AG-groupoid (RC-AG-groupoid) if $\forall a, b, c \in S$,

$$a(bc) = a(cb) \tag{2.2}$$

(c) — a bi-commutative AG-groupoid (BC-AG-groupoid) if it is both LC-AG-groupoid and an RC-AG-groupoid.

Example 1. *We list some examples of smallest order of these AG-groupoids to show their existence. (i) LC-AG-groupoid of order* 3. *(ii) RC-AG-groupoid of order* 3. *(iii) BC-AG-groupoid of order* 3.

*	1	2	3
1	1	1	1
2	1	1	1
3	2	1	1

(i)

*	1	2	3
1	1	1	1
2	1	1	1
3	2	2	1

(ii)

*	1	2	3
1	1	1	1
2	1	1	1
3	2	2	2

(iii)

3. left commutative AG-test

Testing of a groupoid for an AG-groupoid has been well explained by P. V. Protic and N. Stevanovic[1]. In this section we discuss a procedure to test whether an arbitrary AG-groupoid $(G, \cdot)$ is an LC-AG-groupoid or not. For this we define the following binary operations;

$$a \circ b = ab \cdot x \tag{3.1}$$

$$a \star b = ba \cdot x \tag{3.2}$$

Now 2.1 holds if,

$$a \circ b = a \star b \tag{3.3}$$

or

$$a \circ b = b \circ a \tag{3.4}$$

To test whether an arbitrary AG-groupoid is an LC-AG-groupoid, it is necessary and sufficient to check if the operation "$\circ$" and "$\star$" coincide $\forall x \in G$. To this end we check the validity of equation 2.1 or $a \circ b = a \star b$. In other words it is enough to check whether the operation $\circ$ is commutative i.e. $a \circ b = b \circ a$. The tables of the operation "$\circ$" for any fixed $x \in G$ is obtained by multiplying a fixed element $x \in G$ by the elements of the " $\cdot$ " table row-wise. It further gives the tables of the operation "$\star$" if these are symmetric along the main diagonal. Hence it could be easily checked whether an arbitrary AG-groupoid is left commutative AG-groupoid or not.

Example 2. *Check the following AG-groupoid* $(G, \cdot)$ *for an LC-AG-groupoid.*

$\cdot$	1	2	3
1	1	1	1
2	1	1	1
3	2	2	2

Table 1

We can extend Table 1 in the way as described above. It is obvious that the tables constructed for the operation " $\circ$ " on the right of the original table are symmetric about the main diagonal and thus coincide with the " $\star$ " tables as required. Hence $(G, \cdot)$ is an LC-AG-groupoid.

$\cdot$	1	2	3	1			2			3		
1	1	1	1	1	1	1	1	1	1	1	1	1
2	1	1	1	1	1	1	1	1	1	1	1	1
3	2	2	2	1	1	1	1	1	1	1	1	1

4. RIGHT COMMUTATIVE AG-TEST

In this section we discuss a procedure to check an AG-groupoid $(G, \cdot)$ for RC-AG-groupoid, for this we define the following binary operations;

$$a \heartsuit b = a \cdot bx \tag{4.1}$$

$$a \diamondsuit b = a \cdot xb \tag{4.2}$$

(2.2) holds if,

$$a \heartsuit b = a \diamondsuit b \tag{4.3}$$

For any fixed $x \in G$, re-writing x-row of the "$\cdot$" table as an index row of the new table and multiplying it by the elements of the index column to construct table of operation "$\diamondsuit$". These extended tables are given to the right of the original table in the following example.

Similarly the table for the operation "$\heartsuit$" for any fixed $x \in G$ is obtained by taking the elements of x-column of the " $\cdot$ " table as an index row of the new table and multiplying it by the elements of the index column of the original table to construct tables for the operation "$\heartsuit$", which are given downward in the extended table of the following example. If the tables for the operation "$\heartsuit$" and "$\diamondsuit$" coincides for all $x \in G$, then (4.3) holds and the AG-groupoid is left commutative-AG-groupoid in this case.

Example 3. *Check the following AG-groupoid for BC-AG-groupoid.*

$\cdot$	1	2	3
1	1	1	1
2	1	1	1
3	2	2	2

Extend the above table in the way as described above we get the extended form as follow:.

$\cdot$	1	2	3	1	1	1	1	1	1	2	2	2
1	1	1	1	1	1	1	1	1	1	1	1	1
2	1	1	1	1	1	1	1	1	1	1	1	1
3	2	2	1	2	2	2	2	2	2	2	2	2
				1	1	1	1	1	1	1	1	1
				1	1	1	1	1	1	1	1	1
				2	2	2	2	2	2	2	2	2

It is clear from the extended table that the tables for the operations"$\heartsuit$" and "$\diamondsuit$" coincide for every $x \in G$, so $(G, \cdot)$ is an RC-AG-groupoid.

5. ideals in LC-AG-groupoids and RC-AG-groupoids

A subset A of the AG-groupoid S is a left (right) ideal of S if,

$$SA \subseteq A(AS \subseteq A) \tag{5.1}$$

A is a two sided ideal or simply an ideal of S if it is both left and right ideal of S.

Remark 1. [1] *If S is an AG-groupoid and $a \in S$, then by the identity 1.1, it follows that:*

$$(aS)S = \underset{x,y\in S}{\cup}(ax)y = \underset{x,y\in S}{\cup}(yx)a \subseteq Sa.$$

From this we conclude that $(AS)S \subseteq SA$.

Further we have the following remarks for our new classes of AG-groupoids.

Remark 2. *If S is an AG-groupoid with left identity e and $a \in S$, then by the medial property and 2.2, it follows that:*

$$S(aS) = \underset{x,y\in S}{\cup}x(ay) = \underset{x,y\in S}{\cup}(ex)(ay) = \underset{x,y\in S}{\cup}(ea)(xy) \subseteq aS.$$

In general for any $A \subseteq S$ we conclude that $S(AS) \subseteq AS$.

Remark 3. *If S is an LC-AG-groupoid and $a \in S$, then by (1.1 and 2.1), it follows that:*

$$(Sa)S = \underset{x,y\in S}{\cup}(xa)y = \underset{x,y\in S}{\cup}(ax)y = \underset{x,y\in S}{\cup}(yx)a \subseteq Sa.$$

Thus for any $A \subseteq S$ we conclude that $(SA)S \subseteq SA$.

Remark 4. *If S is an RC-AG-groupoid with left identity e and $a \in S$, then by 1.1 and 2.2, it follows that:*

$$S(Sa) = \underset{x,y\in S}{\cup}x(ya) = \underset{x,y\in S}{\cup}(ex)(ya) = \underset{x,y\in S}{\cup}(ex)(ay) = \underset{x,y\in S}{\cup}(ea)(xy) \subseteq aS.$$

Hence in general $S(SA) \subseteq AS$ for $A \subseteq S$.

Remark 5. *If S is an RC-AG-groupoid with left identity e and $a \in S$, then by medial property and 2.1, it follows that:*

$$\begin{aligned}(Sa)S &= \underset{x,y\in S}{\cup}(xa)y = \underset{x,y\in S}{\cup}(xa)(ey) = \underset{x,y\in S}{\cup}(xa)(ye)\\ &= \underset{x,y\in S}{\cup}(xy)(ae) = \underset{x,y\in S}{\cup}(xy)(ea) \subseteq Sa.\end{aligned}$$

Thus $(SA)S \subseteq SA$ for $A \subseteq S$.

Definition 2. [1]*Let S be an AG-groupoid and $A, B \subseteq S$, than A and B are right (left) connected sets if $AS \subseteq B$ and $BS \subseteq A$ ($SA \subseteq B$ & $SB \subseteq A$).*

Remark 6. *If L is a left and R is a right ideal of an LC-AG-groupoid S then by (2.1 and 5.1) and left invertive law, we have*

$$(LR)S = (SR)L = (RS)L \subseteq RL \text{ and } (RL)S = (SL)R \subseteq LR.$$

It follows that LR and RL are right connected sets.

Proposition 1. *Let S be an LC-AG-groupoid then for each $a \in S$ the set $a \cup aS$ and $aS \cup Sa$ are right connected sets.*

Proof. If $a \in S$, then by Remarks (1 and 3), we have $(a \cup Sa)\, S = aS \cup (Sa)S \subseteq aS \cup Sa$, also,

$$(aS \cup Sa)\, S = (aS)S \cup (Sa)S \subseteq Sa \cup Sa \subseteq a \cup Sa.$$

Hence the result follows. ∎

Proposition 2. [1] *Let S be an AG-groupoid and let A and B are right connected sets, then $A \cap B$(if $A \cap B \neq \emptyset$) and $A \cup B$ are right ideals of S.*

Proof. We have,

$$(A \cap B)\, S \subseteq AS \cap BS \subseteq B \cap A$$
$$(A \cup B)\, S = AS \cup BS \subseteq B \cup A.$$

Hence the result follows. ∎

Theorem 1. *Let S be an LC-AG-groupoid then for each $a \in S$ the set $a \cup aS \cup Sa$ is right ideal of S.*

Proof. Let $a \in S$, then by Remarks (1-3), we have

$$(a \cup aS \cup Sa)\, S = aS \cup (aS)S \cup (Sa)S \subseteq aS \cup Sa \cup Sa \subseteq aS \cup Sa \subseteq a \cup aS \cup Sa.$$

Hence the result follows. ∎

Theorem 2. *Let S be an LC-AG-groupoid then for each $a \in S$ the set $R(a) = (aS)a$ is the minimal right ideal of S containing a.*

Proof. Let $a \in S$, then by medial property and (1.1 and 2.1), we have

$$((aS)a)\, S = \underset{x,y \in S}{\cup}((ax)a)y = \underset{x,y \in S}{\cup}(ya)(ax) = \underset{x,y \in S}{\cup}(ay)(ax) =$$
$$= \underset{x,y \in S}{\cup}(aa)(yx) = \underset{x,y \in S}{\cup}(yx \cdot a)a = \underset{x,y \in S}{\cup}(a \cdot yx)a = (aS)a.$$

Hence, $R(a)$ is a right ideal of S. Now, let R be a right ideal of S such that $a \in R$. Then we have

$$R(a) = (aS)a \subseteq (RS)R \subseteq RR \subseteq R.$$

So $R(a)$ is a minimal right ideal of S containing a. In general, if $A \subseteq S$. Then $(AS)A$ is a minimal right ideal generated by A. ∎

Theorem 3. *Let S be an RC-AG-groupoid with left identity e then for each $a \in S$ the set $J(a) = a \cup aS \cup Sa$ is the minimal (two sided) ideal of S containing a.*

Proof. By Proposition 2 and Remarks (2 and 4), we have

$$S\,(a \cup aS \cup Sa) = Sa \cup S(aS) \cup S(Sa) \subseteq Sa \cup aS \cup aS \subseteq Sa \cup aS \subseteq a \cup aS \cup Sa.$$

Thus $J(a)$ is a left ideal. Now again by Proposition 2 and Remarks (1 and 3), we have$(a \cup aS \cup Sa)\, S = aS \cup (aS)S \cup (Sa)S \subseteq aS \cup Sa \cup Sa \subseteq aS \cup Sa \subseteq a \cup aS \cup Sa$. Thus $J(a)$ is a right ideal, and hence it is a two sided ideal or simply an ideal of S. If J is an ideal of S and $a \in J$ then

$$J(a) = a \cup (aS \cup Sa) \subseteq J \cup (JS \cup SJ) \subseteq J \cup (J \cup J) \subseteq J \Rightarrow J(a) \subseteq J.$$

Hence the result follows. ∎

Theorem 4. *If S is an RC-AG-groupoid with left identity e then for $a \in S$, the sets $a(Sa)$ and $(aS)a$ are ideals of S. If $a \in a(Sa)$ (resp. $a \in (aS)a$) then $a(Sa)$ (resp. $(aS)a$) is a minimal ideal generated by a. Further if $a \in a(Sa) \cap (aS)a$, then $(aS)a = a(Sa)$ and it is minimal ideal generated by a.*

Proof. If $a \in S$, then by the paramedial property (1.1) and 2.2, we have

$$\begin{aligned} S\,(a(Sa)) &= \underset{x,y\in S}{\cup} x(a(ya)) = \underset{x,y\in S}{\cup}(ex)(a\cdot ya) = \underset{x,y\in S}{\cup}(ea)(x\cdot ya) = \\ &= \underset{x,y\in S}{\cup} a(x\cdot ay) = \underset{x,y\in S}{\cup} a(ay\cdot x) = \underset{x,y\in S}{\cup} a(xy\cdot a) \subseteq a(Sa). \end{aligned}$$

Similarly,

$$\begin{aligned} (a(Sa))S &= \underset{x,y\in S}{\cup}(a(xa))y = \underset{x,y\in S}{\cup}(y\cdot xa)a = \underset{x,y\in S}{\cup}(y\cdot ax)a = \\ &= \underset{x,y\in S}{\cup}(y\cdot ax)(ea) = \underset{x,y\in S}{\cup}(y\cdot ax)(ae) = \underset{x,y\in S}{\cup}(e\cdot ax)(ay) = \\ &= \underset{x,y\in S}{\cup}(ea)(ax\cdot y) = \underset{x,y\in S}{\cup} a(yx\cdot a) \subseteq a(Sa) \end{aligned}$$

Hence, $a(Sa)$ is an ideal of S.

Now, again using the paramedial law, and (1.1) and 2.2, we have

$$\begin{aligned} S\,((aS)a) &= \underset{x,y\in S}{\cup} x(ay\cdot a) = \underset{x,y\in S}{\cup}(ex)(ay\cdot a) = \underset{x,y\in S}{\cup}(e\cdot ay)(xa) = \\ &= \underset{x,y\in S}{\cup}(e\cdot ay)(ax) = \underset{x,y\in S}{\cup}(x\cdot ay)(ae) = \underset{x,y\in S}{\cup}(x\cdot ay)(ea) = \\ &= \underset{x,y\in S}{\cup}(ex\cdot ay)(a) = \underset{x,y\in S}{\cup}(ea\cdot xy)a = \underset{x,y\in S}{\cup}(a\cdot xy)a \subseteq (aS)a \end{aligned}$$

Similarly,

$$\begin{aligned} S\,((aS)a) &= \underset{x,y\in S}{\cup}((ax\cdot a)y) = \underset{x,y\in S}{\cup} ya\cdot ax = \underset{x,y\in S}{\cup} yx\cdot aa = \\ &= \underset{x,y\in S}{\cup}(e\cdot yx)(aa) = \underset{x,y\in S}{\cup}(a\cdot yx)(ae) = \underset{x,y\in S}{\cup}(a\cdot yx)(ea) \subseteq (aS)a. \end{aligned}$$

Hence $(aS)a$ and $a(Sa)$ are ideals of S. If A is an ideal on S, then for every $a \in A$ we have $(aS)a \subseteq A$ and $a(Sa) \subseteq A$, clearly. If $a \in A \cap (aS)a$ (resp. $a \in A \cap a(Sa)$), then $(aS)a$ (resp. $a(Sa)$) is a minimal ideal generated by a. If $a \in A \cap (aS)a \cap a(Sa)$, then by minimality, it follows that $(aS)a = a(Sa)$. Clearly, for each $a \in S$ it holds that $(aS)a \subseteq Sa$ and $a(Sa) \subseteq Sa$. ∎

6. characterization of BC-AG-groupoid

In this section we discuss the relations of BC-AG-groupoid with already known classes of AG-groupoids. We start with the following results which proves that every AG*-groupoid is RC-AG-groupoid, but not an LC-AG-groupoid as in Example 4.

Theorem 5. *Every AG*-groupoid is RC-AG-groupoid.*

Proof. Let S be an AG*-groupoid, and $a, b, c \in S$. Then $a(bc) = (ba)c = (ca)b = a(cb) \Rightarrow a(bc) = a(cb)$. Hence S is RC-AG-groupoid. ∎

The following example shows that every AG*-groupoid is not an LC-AG-groupoid.

Example 4. *Let $S = \{1, 2, 3, 4, 5, 6\}$ then it is easy to verify that S is an AG*-groupoid, but not an LC-AG-groupoid as, $(1 * 2) * 1 \neq (2 * 1) * 1$.*

*	1	2	3	4	5	6
1	3	4	5	5	5	5
2	3	4	6	6	5	5
3	5	5	5	5	5	5
4	6	6	5	5	5	5
5	5	5	5	5	5	5
6	5	5	5	5	5	5

Now we prove the following:

Theorem 6. *Every LC-AG*-groupoid is a semigroup.*

Proof. Let S be an LC-AG-groupoid satisfying AG*-groupoid property, and $a, b, c \in S$. Then $(ab)c = (ba)c = a(bc) \Rightarrow (ab)c = a(bc)$. Hence S is a semigroup. ∎

Theorem 7. *Every BC-AG-groupoid is paramedial AG-groupoid.*

Proof. Let S be a BC-AG-groupoid, and $a, b, c, d \in S$. Then $ab \cdot cd = ab \cdot dc = ba \cdot dc = bd \cdot ac = db \cdot ac = db \cdot ca \Rightarrow ab \cdot cd = db \cdot ca$. Hence S is paramedial AG-groupoid. ∎

It follows that:

Corollary 1. *Every BC-AG-groupoid is left nuclear square AG-groupoid.*

The following counterexamples shows that neither AG**-groupoid nor BC-AG-groupoid is a nuclear Square AG-groupoid. However, both these properties jointly gives the desired relation as given in the following theorem.

Example 5. *(i). AG**-groupoid of order 3 that is not a nuclear square AG-groupoid. (ii). BC-AG-groupoid that is not nuclear square AG-groupoid.*

·	1	2	3
1	1	1	1
2	1	1	1
3	1	2	2

(i)

·	1	2	3
1	1	1	1
2	1	1	1
3	2	2	1

(ii)

Theorem 8. *Let S be a BC-AG**-groupoid. Then the following conditions are equivalent.*

(i) S is middle nuclear square AG-groupoid;
(ii) S is right nuclear square AG-groupoid;
(iii) S is nuclear square AG-groupoid.

Proof. Let S be a BC-AG**-groupoid. Then

$(i) \Rightarrow (iii)$. Assume (i) holds, let $a, b, c \in S$. Then $a(bc^2) = b(ac^2) = b(c^2a) = (bc^2)a = (c^2b)a = (ab)c^2 \Rightarrow a(bc^2) = (ab)c^2$. Thus S is right nuclear square AG-groupoid. Now, by lemma (1) S is left nuclear square AG-groupoid as well. Hence S is nuclear square AG-groupoid which is (iii).

$(iii) \Rightarrow (ii)$ Obvious. Finally we show,

$(ii) \Rightarrow (i)$. Assume (ii) holds, and let $a, b, c \in S$. Then $a(b^2c) = b^2(ac) = b^2(ca) = c(b^2a) = c(ab^2) = (ca)b^2 = (b^2a)c = (ab^2)c \Rightarrow a(b^2c) = (ab^2)c$. Which proves (i).

Hence the theorem is proved. ∎

Now we give an example of left alternative AG-groupoid and BC-AG-groupoid that are not flexible AG-groupoid.

Example 6. *(i). left alternative AG-groupoid, which is not flexible AG-groupoid. (ii). BC- AG-groupoid, which is not flexible AG-groupoid.*

*	1	2	3	4
1	3	3	3	2
2	4	3	3	3
3	3	3	3	3
4	3	1	3	3

(i)

*	1	2	3
1	2	2	2
2	3	3	3
3	3	3	3

(ii)

However, we have the following:

Theorem 9. *Every left alternative BC-AG-groupoid is flexible AG-groupoid.*

Proof. Let S be a BC-AG-groupoid satisfying also the left alternative AG-groupoid property, and let $a, b \in S$. Then $ab{\cdot}a = ba{\cdot}a = aa{\cdot}b = a{\cdot}ab = a{\cdot}ba \Rightarrow ab{\cdot}a = a{\cdot}ba$. Hence S is flexible AG-groupoid. ∎

Theorem 10. *Every LC-AG-groupoid with left identity e is commutative semigroup.*

Proof. Let S be a RC-AG-groupoid with left identity e, and let $a, b \in S$. Then $ab = ea \cdot eb = ae \cdot eb = (eb \cdot e)a = (be \cdot e)a = (ee \cdot b)a = ba \Rightarrow ab = ba$. Hence$S$ is commutative, but commutativity implies associativity in AG-groupoids, hence S is commutative semigroup. ∎

One can easily verify that neither T^1-AG-groupoid nor BC-AG-groupoid is self dual AG-groupoid.

Example 7. *(i). T^1-AG-groupoid of order 3 which is self dual AG-groupoid. (ii). BC- AG-groupoid of order 3 which is not self dual AG-groupoid.*

·	1	2	3
1	1	1	1
2	1	1	3
3	1	2	1

(i)

·	1	2	3
1	1	1	1
2	1	1	1
3	2	2	2

(ii)

However, we prove the following:

Theorem 11. *Every T^1-BC-AG-groupoid is self dual AG-groupoid.*

Proof. Let S be a T^1-BC-AG-groupoid, and let $a, b, c \in S$. Then $a(bc) = a(cb)$ $\Rightarrow (bc)a = (cb)a = (ab)c = (ba)c \Rightarrow (bc)a = (ba)c \Rightarrow a(bc) = c(ba)$. Hence S is self dual AG-groupoid. ∎

Theorem 12. *Every BC-AG-3-band is commutative semigroup.*

Proof. Let S be BC-AG-3-band, and let $a, b \in S$. Then $ab = (a \cdot aa)(b \cdot bb) = (ab)(aa{\cdot}bb) = (ba)(aa{\cdot}bb) = (ba)(ab{\cdot}ab) = (ba)(ba{\cdot}ab) = (ba)(ba{\cdot}ba) = (ba)(bb{\cdot}aa)$ $= (b{\cdot}bb)(a{\cdot}aa) = ba \Rightarrow ab = ba$. Thus S is commutative and hence is associative. Equivalently S is commutative semigroup. ∎

An element a of an AG-groupoid S is called left cancellative if $ab = ac \Rightarrow b = c$, right cancellative and cancellative elements are defined analogously.

Theorem 13. *Every LC-AG-groupoid with a cancellative element is a commutative semigroup.*

Proof. Let S be an LC-AG-groupoid with a cancellative element x, and let $a, b \in S$. Then $ab \cdot x = ba \cdot x \Rightarrow ab = ba$. Thus S is commutative, but commutativity implies associativity in AG-groupoids. Hence S commutative semigroup. ∎

References

[1] P.V. Protic and S. Nebojsa, AG-test and some general properties of AG-groupoids, PU.M.A. Vol. 6(4), 371-383, (1995).
[2] GAP: Groups Algorithm and Programming, Version 4.4.12, 2008, (2012).
[3] A. Distler, Shah M. and V. Sorge, Enumeration of AG-groupoids, Lecture Notes in Computer Science, Volume 6824/2011, 1-14 (2011)
[4] M. Rashad, I. Ahmad, Amanullah and M. Shah, On relations between right alternative and nuclear square AG-groupoids, Int. Math. Forum, Vol. 8(5), 237-243, 2013.
[5] I. Ahmad, M. Rashad and M. Shah, Some Properties of AG*-groupoid, Res. J. Recent Sci.,Vol. 2(4), 91-93, April (2013)
[6] I. Ahmad, M. Rashad and M. Shah, Some new result on T^1, T^2 and T^4-AG-groupoids, Research Journal of Recent Sciences, Vol. 2(3), 64-66, (2013).
[7] M. Rashad, I. Ahmad, and M. Shah, Left transitive AG-groupoids, arxiv.org.
[8] M. Shah, A theoretical and computational investigation of AG-groups, PhD thesis, Quaid-i-Azam University Islamabad, (2012).
[9] M. Rashad, I. Ahmad, and M. Shah, Stein AG-groupoids, arxiv.org
[10] J. R. Cho, J. Jezek and T. Kepka, Paramedial Groupoids, Czechoslovak Mathematical Journal, 49 (124) 1996.
[11] M. Shah, I. Ahmad and A. Ali, Discovery of new classes of AG-groupoids, Res. J. Recent Sciences, Vol, 1(11), 47-49 (2012).
[12] M. Khan, Faisal, V. Amjad, On some classes of Abel-Grassmann's groupoids. J. Adv. Res. Pure Math. 3(4), 109–119 (2011)
[13] M. Shah, I. Ahmad and A. Ali,, On introduction of new classes of AG-groupoids, Res. J. Recent Sciences, Vol. 2(1), 67-70, January (2013).
[14] M. Rashad, I. Ahmad, and M. Shah, Some general properties of LAD and RAD-AG-groupoids, www.arXiv.org 1403.5218.
[15] Amanullah, M. Rashad, I. Ahmad, M. Shah, On Modulo AG-groupoids, arXiv: 1403.2564
[16] Amanullah, M. Rashad, I. Ahmad, M. Shah, M. Yousaf, Modulo Matrix AG-groupoids and Modulo AG-groups arXiv:1403.2304

E-mail address: rashad@uom.edu.pk

E-mail address: iahmaad@hotmail.com

A. Department of Mathematics, University of Malakand, Chakdara Dir(L), Pakistan.

E-mail address: shahmaths_problem@hotmail.com

B. Department of Mathematics, Government Post Graduate College Mardan, Pakistan.